### **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification		(11) International Publication Number: WO 88/09367
C11D 3/386	A1	(43) International Publication Date: 1 December 1988 (01.12.88)
(21) International Application Number: (22) International Filing Date: 31	PCT/US88/018	(75) Inventors/Applicants (for US only): KOLATTUKUDY,
(31) Priority Application Number: (32) Priority Date: 29	056,5 May 1987 (29.05.8	(74) Agent: PASSE, James, G.; Genencor, Inc., 180 Kimball
(33) Priority Country:  (60) Parent Application or Grant (63) Related by Continuation US Filed on 29	056,500 (Cl May 1987 (29.05.8	
(71) Applicant (for all designated State ENCOR, INC. [US/US]; 180 Ki Francisco, CA 94080 (US).	es except US): GE mball Way, So. S	Published  With international search report.  Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.
		*

(54) Title: CUTINASE CLEANING COMPOSITION

### (57) Abstract

This invention relates to cleaning compositions and methods for using them. Particularly, the invention relates to compositions comprising a surfactant and a cutinase enzyme.

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

					•
AT	Austria:	FR	France	ML	Mali
ΑŪ	Australia:	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	ľT	Italy	NO	Norway
BJ	Benin	JР	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG.	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland ·	LÍ	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
DE	Germany, Federal Republic of	LU	Luxembourg	TG	Togo
DK	Denmark	MC	Monaco	US	United States of America
F	Finland	MG	Madagascar		ombe omes of America

### Cutinase Cleaning Compositions

### Background of the Invention

### a) Field of the Invention

The invention relates to enzymatic cleaning compositions and methods for using them. Particularly the invention relates to cleaning compositions comprising a surfactant and a cutinase enzyme.

### b) Background Information

A wide variety of enzymes are well known for use in cleaning compositions. The use of B. subtilisins and B. licheniformis protease in commercial preparations is common. Other enzymes have also been used in commercial cleaning compositions such as, for example, U.S. Patent No. 4,011,169, and British Patnet No. 1,293,613. Also a comprehensive review article of lipases in cleaning compositions can be found in Journal of Applied

Biochemistry, 2:218-229 (1980) in an article entitled "Lipases as Detergent Components". Lipolytic detergent additives are also known from e.g., British Patent Specification No. 1,293,613 and Canadian Patent No. 835,343.

U.S. Patent No. 3,950,277 and British Patent Specification No. 1,442,418 disclose lipase enzymes combined with an activator and calcium and/or magnesium ions, respectively, which are utilized to pre-soak soiled fabrics and to remove triglyceride stains and soils from polyester or polyester/cotton fabric blends, respectively, Suitable microbial lipases for use therein (apart from animal and plant derived lipases) are said to be those derived from <u>Pseudomonas</u>, <u>Aspergillus</u>, <u>Pneumococcus</u>, <u>Staphylococcus</u>, and <u>Staphylococcus</u> toxins, <u>Mycobacterium tuberculosis</u>, <u>Mycotorula lipolytica</u>, and <u>Sclerotinia</u>.

British Patent Specification No. 1,372,034 discloses a detergent composition comprising a bacterial lipase produced by <u>Pseudomonas stutzeri</u> strain ATCC 19154. Furthermore, it is recommended that the preferred lipolytic enzymes should have a pH optimum between 6 and 10, and should be active in said range, preferably between 7 and 9. Around 1970, this presumed <u>Pseudomonas stutzeri</u> strain was reclassified as <u>Pseudomonas aeruginosa</u>, as appears for example from the ATCC catalogues.

European Patent Application EP-A-0130064 discloses an enzymatic detergent additive comprising a lipase isolated from <u>Fusarium</u> oxysporum with an alleged higher lipolytic cleaning efficiency than conventional lipases.

In European Patent Application No. 0214761, Enzymatic detergent additives are described as the active component of which is a microbially produced lipase from a strain of <u>Pseudomonas cepacia</u>. The lipases described therein are claimed to be superior to the lipolytic detergent action of the prior art, especially at low temperature washing processes (around 60°C and below).

In PCT Patent Application No. 87/00859 other novel lipolytic enzymes are described as having an optimal pH in the range of 8 - 10.5 at a temperature of 60°C or less from bacterial strains selected from <u>Pseudomonas pseudoalcaligenes</u>, <u>P. stutzeri</u> and <u>Acinetobacter calcoaceticus</u>. These enzymes are described as particularly effective at low temperatures; i.e., 40°C or lower and effective in both liquid and solid detergent compositions.

Also in U.S. Patent No. 3,950,277, it is described in general terms that lipases from Pseudomonas are suited as agents for removal of oily stains from fabrics, if used together with a special group of lipase activators. The art cited does not, however, cover cutinase enzymes from Pseudomonas or any other microbial source. However, prior art enzymes for use in cleaning compositions, while effective on many proteins and lipids, are not completely effective against all stains commonly found in laundry and other cleaning applications.

Further, many lipases are not stable at pH 8 - 11 where most cleaning compositions are used. Even further, most enzymes for use in cleaning compositions are not very stable, if at all, under oxidative conditions or in the presence of other enzymes such as proteases.

### SUMMARY OF THE INVENTION

Accordingly it has been discovered that combinations of a surfactant and a substantially pure microbial cutinase enzyme are effective compositions for cleaning applications. The cutinase enzyme preparations possess activity at pH of from about 8 and 11, exhibit cleaning activity in aqueous solution at concentrations from about .05 mg/L to about 100 mg/L or more at temperatures from about 20°C to about 50°.

The enzymes are oxidatively stable and stable in the presence of other enzymes such as proteases. Even further, the cutinases show a synergistic effect when a plurality of surfactants are used with the cutinase.

The invention also relates to the improved process for enzymatically cleaning a material with an aqueous solution; the improvement comprising adding a substantially pure cutinase to the cleaning solution.

#### DETAILED DESCRIPTION OF THE INVENTION

Applicant has discovered that cutinase enzymes are useful when included in cleaning compositions. These compositions may take on a variety of forms such as for laundry cleaning, household and industrial cleaning, and the like. The cleaning compositions comprise combinations of known surfactants and a microbial cutinase enzyme which can be used to clean a wide variety of materials. The composition can be added to aqueous solution or solid powder, or formulated in an aqueous solution or solid powder and used according to conventional cleaning techniques.

Enzyme

Cutinases are well known in the art and are available from a wide variety of sources. See Cutinases from Fungi and Pollen, P.E. Kolattukudy, pg. 472 - 504, incorporated herein by reference, for discussion of cutinases useful in the practice of the invention. A preferred cutinase is that cutinase isolated in a substantially pure form from Pseudomonas putida, particularly the P. putida, ATCC 53552, described in copending U.S. patent application, Serial No. 932,959 filed November 19, 1986 and incorporated herein by reference, which enzyme therefrom has the following amino acid sequence:

0936	7								-		PCT
	•			•		5			. 10		
1414	010	lev :	<b>p</b> r•	410	th/	pro	• • •	6'6'		<b>p</b> h	e pro
414	v41	414	418	phe	419	419	10	• •	<b>)</b> ′(	ty	thr
ihr	ser	ser	gla	140	30	894	910	10.	cys	419	11e
tye	1479	<b>310</b>	40	419	lev	91y	· ·••••	1'1	gly	ya 1	arg
-	50	•						١,			60
his	pro	*41	He	lev	tre	117	418	1'1	thr	gly.	414
gly	<b>)</b> r•	ser	the	tyr	ata	gly	leu	le.	. 30 ser	Als	tṛp
414	ser	his	gly	phe	v41	<u>.</u> 141	414	414	414	970	thr
Ser	410	a1a	gly	thr	90 . 917	4/1	910	E4:	Tev	ala	cys
leu	419	tyr	100 1eu	****	14	glv	410	<b>013</b>	thr	)re	tyr
117	110	tyr	ser	gly	lys	tev	414	ter	gly	arg	120 741
917	thr	307	gly .	his.	. ser	g 1 n	gly	1. 1813	130	917	ser
							140				
130	ert	<b>a 1,a</b>	#1y	e î a	459	thr.	418	49,	· Arg	thr	the
a1a	pr•	11e	gla	; pre	150	thr	lev	91,	· lev	gly	,his
45)	ser	414	160 ser	gla	arg	arg	gla	gla	gly	<b>)</b> re	set
ghé	170 1eu	net	ser	917	914	gly	459	tar	11e	414	160 phe
<b>pro</b>	tyr	lew	414	ala	gla	910	<b>va1</b> -	tyr	190	arg	41a-
45R	val	<b>970</b>	val	phe	trp	917	\$00 \$1.	4/5	479	tyr	val
ser-	Als	phe	glu	<b>»</b> ••	210 741	917	ser	<b>9</b> 1y	817	414	tye
arg	917	<b>970</b>	250 550	the	ala	trp	pre	arg	phe	gla	leu
art	230 419	410	gla	459	ala	arg	414	thr	Phe	tyr	240 11y
			440	١				10.	250		•

Other sources of bacterial and fungal cutinases include:

Fusarium solani pisi
Fusarium roseum sambucinum
Fusarium roseum culmorum
Helminthosporum sativum
Ulocladium consortiale
Streptomyces scabies
Colletotrichum capsici
Phytopthora cactorum
Botrytis cineria
Colletotrichum gloeosporioides

The cutinase of the invention should preferably be selected to cause at least about 10%, and preferably 20%, hydrolysis of the given fat under given conditions. Normally the amount would be in a concentration of from about .01% to about 5.0% by weight of the surfactant, and preferably from about .05% to about 3%, such that upon dilution in wash water it is in a concentration of at least about .05mg/L. Further, one skilled in the art could take the preferred cutinase or, for that matter, any cutinase of the invention or any immuniologically identical cutinase and use random or selective replacement of amino acids to produce other cutinases which are more or less selective toward given substrates or include modification in activity such as oxidative stability. Substantially pure cutinase includes the isolated enzyme as well as the broth containing the enzyme in unpurified form but essentially free of other enzymes and enzyme sources.

The natural substrate of cutinase is cutin which is a biopolyester polymer which covers the plant leaves, fruits, etc., see Structure, Biosynthesis and Biodegradation of Cutin and Suberin. (1981), P.E. Kolattukudy, Arm. Rev. Plant Physiol., 32, pgs. 539-567. Stains comprising lipids which could be hydrolyzed or bound by cutinase on a substrate such as cloth would be similar to the natural substrate cutin. Cutinase, for these types of stains, will be more

effective than the prior art lipases. The cutinases will work especially well on gravy, oils and greases, plant or grass, oil based makeup and collar stains.

Cutinases are distinguishable from other lipases by methods well known in the art, See R.E. Purdy and P.E. Kolattukudy, <u>Biochemistry</u>, "Cutinase Assay", 14:2831-2840, (1975). Microbial cutinases from both fungal and bacterial sources have very good activity at pH 8 - pH 11 which is an ideal pH condition for detergent use.

Because of the specific activity of cutinases, it is a preferred aspect of the invention to combine one or more other cutinases, or one or more other enzymes, such as proteases, amylases or other lipases, along with the cutinase of the invention in the cleaning composition. Further, Applicant shows a synergistic increase in hydrolytic activity of cutinase when two or more surfactants are combined along with the cutinase enzyme.

Cutinases then are ideal for cleaning composition inclusion. They have stability oxidatively such as in  $H_2O_2$ . They have good stability in a temperature range of from about  $20-50^{\circ}\text{C}$  which is ideal from a cleaning point of view. They are also stable in the presence of other enzymes; e.g., proteases, and as such, are ideal for mixtures of enzymes.

### The Surfactant

A number of known compounds are suitable surfactants useful in the present compositions. These include nonionic, anionic, cationic, or zwitterionic detergents, as disclosed in U.S. 4,404,128 to Barry J. Anderson and U.S. 4,261,868 to Jiri Hora et al. The art is familiar with the different formulations which can be used as cleaning compositions.

# The Cleaning Compositions and Method of Use

Outinases can be formulated as a purposefully added ingredient into known powdered and liquid detergents having pH between 6.5 and 12.0 at levels of about 0.01 to about 5% (preferably .05 to .5%) by weight of the detergent. These detergent cleaning compositions can also include other enzymes such as known proteases and amylases, as well as bleaches, colorants, builders, and stabilizers.

The cutinase of the invention may be added to powdered detergents in the form of granulates or prills, prepared by methods known in the art such as described in British Patent Nos. 1,324,116 and 1,362,365 and U.S. patent Nos. 3,519,570; 4,106,991 and 4,242,219.

The cutinase preparations of the invention can be prepared by cultivating the microorganisms defined herein or otherwise cutinase containing microorganism under appropriate conditions. In order to obtain reasonable yields of enzyme, media containing readily assimilable carbon and energy sources as necessary such as a nitrogen source, as well as calcium and magnesium salts and trace elements and cutin, or monomers of cutin, or compounds resembling cutin or cutin monomers. One could also obtain the gene for cutinan and express in any organism of choice where one may not have to add cutin or cutin monomers into the fermentation.

The addition of cutinase to conventional cleaning compositions does not create any special use limitation. In other words, any temperature and pH suitable for detergent compositions containing enzymes is also suitable for the present compositions.

Although the preferred form of the invention has been described above, it will be obvious to those skilled in the art to which the invention pertains, that, after understanding the invention and in view of the following testing as a whole, various changes and equivalent modifications may be made without parting from the scope of the invention as defined by the claims.

9

### STABILITY CUTINASE AGAINST PROTEASES

Reaction conditions:

Buffer: 0.1 M NaP, pH 10

Temp: 37°C

Lipase: 42 ug/ml

Approximately 1:1 protease: cutinase aqueous solution were made up with the following results.

#### ENZYME ACTIVITY

#### **Protease**

Incubation Time	o min	5 min	10 min	15 min	14 hrs
None	1.57	1.60	1.47	1.63	1.61
Maxacal (35ug/ml)	1.68	1.58	1.72	1.66	0.134
Esperase	1.73	1.64	1.59	1.51	0.456
(64 ug/ml)					

Maxical is Gist-Brocade's brand of subtilisin enzyme (protease)

Esperase is Novo's brand of protease enzyme (protease)

# TEMPERATURE STABILITY OF BACTERIAL CUTINASE

### HALF LIFE AT 50°C

<u>Hq</u>	Hrs.
7	30
8	25
9	12
10	0.3

Enzyme was incubated at 50°C in 0.1M sodium phosphate buffer at various pH's and activity was measured by hydrolysis of trioctanoin in polyvinyl alcohol emulsions.

# EFFECT OF DETERGENTS ON HYDROLASE ACTIVITY

### Reaction Conditions:

substrate: p-nitro-phenyl butyrate, lmm (pmp)

pH: 8.0

buffer: 0.lm tris pH 8.0

temperature: 25°C

enzyme: bacterial cutinase from ATCC 53552

An aqueous solution with the following were made up and the enzyme activity was measured in these solutions using pmp as a substrate by following absorbance of p-nitrophenol at 410 mm.

Triton x100%	SDS &	% Activity
. •		
0	. 0	100
0.2		78
0.4		60
· <del></del>	0.05	30
	0.1	23
	0.2	14
	0.4	6
0.4	0.4	78
0.2	0.2	98
0.2	0.05	125
0.2	0.1	138

6.1 0.1 130 0.05 0.05 132


- 1) Non-ionic detergent inhibition is not significant at low concentrations.
- 2) An ionic detergent inhibitor at high concentrations.
- 3) Mixture of anionic and non-ionic detergents stimulate activity.

# STABILITY TOWARD OXIDANTS


Cutinase .5 mg/ml in 0.1m sodium phosphate buffer, was incubated with various levels of hydrogen peroxide at pH 8.4 and 25°C for 2 hours, and hydrolytic activity was measured by a pH-stat using trioctanoim-polyvinyl alcohol emulsion.

[H <sub>2</sub> 0 <sub>2</sub> ppm]	Hydrolytic Activity	•
	% Remaining	
0	100	
100	86	
200	86	
500	91	
1000	95	

Act. (mmol/mg.mln)



Act. (mmol/mg.min)



SUBSTITUTE SHEET

### What is Claimed is:

- 1. An enzymatic cleaning composition comprising a substantially pure microbial cutinase in an amount sufficient that upon dilution in an aqueous solution for use there is sufficient cutinase to achieve a concentration of from about .05 mg/L to about 100 mg/L of cutinase.
- 2. A composition according to Claim 1 wherein the cutinase is the cutinase present in an amount of from about 0.01% to about 5% by weight of the surfactant.
- 3. A composition according to Claim 2 wherein the <u>Pseudomonas</u> putida is ATCC 53552.
- 4. A composition according to Claim 1 which further comprises one or more enzymes selected from lipases, amylases or proteases.
- 5. A composition according to Claim 1 wherein the cutinase selected is capable of causing at least 10% hydrolysis of a cutin or cutin-like substrate desirous of being cleaned from a given surface.
- 6. A composition according to Claim 1 wherein the cutinase is a non-naturally occurring cutinase having at least one amino acid randomly or selectively replaced by an amino acid not naturally found at that position.

- A composition according to Claim 1 wherein the cutinase is added in the form of cell fermentation broth containing cutinase but essentially free of other enzymes or enzyme sources.
- An enzymatic cleaning additive wherein the active component 8. which is a substantially pure microbially produced cutinase.
- A composition according to Claim 8 which further comprises one 9. or more enzymes selected from lipases, amylases or proteases.
- 10. A composition according to Claim 8 comprising a plurality of surfactants.
- A composition according to Claim 10 wherein the surfactants are SDS and Triton X-100.
- 12. An improved method for enzymatically cleaning a material having a cutin or cutin-lke stain comprising:
  - a) selecting a cutinase enzyme;
  - b) forming an aqueous solution with said enzyme in a concentration of from about 0.05 mg/L to about 100 mg/L;
  - c) contacting the material with the solution of step b); and
  - d) rinsing the material of step c).
- 13. A method according to Claim 12 wherein the aqueous solution further comprises a surfactant compatible with the selected cutinase.

18
14. A method according to Claim 13 wherein the aqueous solution further comprises a surfactant compatible with the selected cutinase.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 88/01844

	FICATIO	OF SUBJECT M	ATTER (if several classi	location symbol	tion and IPC	ate div	
According	to Internati	anal Patent Classific	ation (IPC) or to both Nat	DUST CHESSINGS	2011 Z.110 1.1 C		•
PC4:	С	11 D 3/38	56	•	·		
CIEL DS	SEARCH	ED					
. PIELDS	302.1.0	<del></del>	Minimum Documen	ntation Searche	d 7	<u> </u>	
assificatio	n System	·		Classification :	Symbols		
			•				
PC4		C 11	D; C 12 N				
•	. [		•		· .	<del></del>	
		Docui to the E	mentation Searched other stent that such Documents	then Minimum I are Included I	Documentation the Fields S	n earched <sup>8</sup>	
•				••			
4							
II. DOCU	MENTS C	ONSIDERED TO	BE RELEVANT	consists of the	relevant pass	ages 12	Relevant to Claim No. 13
tegory •	Citati	on of Document, 13	with indication, where app	ropriate, or an			
P,X	EP,	see exam	152 (GENENCON ple 13; abst	R) 25 Ma ract; 1	ay 1988 page 1,	3	1-3,5,6,8, 12,13
ĺ	cit	lines 29 ed in the	9-30 e application	ı	•		
							* .
	•						
	•					*	
	•						
							_
			••		•		
			•				
	٠					•	
į		• •			• •		
l		•			•	·	j
			8.	•			
				•			
"A" doc con "E" eari filin "L" doc white cita "O" doc oth "P" doc late	ument defi- sidered to ier docume g date ument whi ch is cited tion or oth ument refe er means ument pub r than the	on the published on the may throw doub to establish the pu- er special reason (a rring to an oral disc lished prior to the in priority date claimed	te of the art which is not vance or after the international its on priority claim(s) or blication date of another a specified) closure, use, exhibition or ternational filling date but	or pricited inventions of the cannot be cannot	lority date an to understan tion ment of part or be consider or an inventive ment of part or be consider ment is comb a, such comb a art.	of the principicular relevanted novel of estep icular relevanted to involve inned with one ination being rof the same	
20+h	Augus	st 1988				4	SET 1300
29 (11				Signature	A American	Officer	

# ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

UŞ 8801844

SA 22728

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 08/09/88

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report		Patent document Publication cited in search report date		Patent family member(s)	Publication date		
	EP-A-	0268452	25-05-88	AU-A- 8115387 AU-A- 7901387	26-05-88 09-06-88		
	•				. •		
			.*				
			•	·*	•		
	•						
	•	• • •	*				